



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NEW BIRD FOR OHIO.

RED-LEGGED BLACK DUCK—(*Anas obscura rubripes*.)

BY REV. W. F. HENNINGER.

THE recent article of Mr. Wm. Brewster in the *Auk*, (Vol. XIX., April, 1902, pp. 183–188), and his foot-note that probably a large proportion of the autumn birds in the Mississippi Valley and about the Great Lakes belong to the form *rubripes* of the Black Duck, brought back to my memory the fact that I received on March 27, 1901, a very large Black Duck, in which I noticed the very bright red feet and yellow bill, and my notebook still contains the remark concerning this specimen, "tarsus red." On examining the dried skin the feet still show an orange color, the pileum and nape are typical *rubripes* and the spots on the throat and down the sides of the neck are almost typical *rubripes*. This shows up still more, when comparing the specimen, a male, with a typical specimen in my collection (shot March 25, 1899, sex unknown), but the head has a more grayish tinge, the throat almost an orange color.

No. 204 (coll. W. F. H.), March 25, 1899, typical *obscura*, measures: wing 24.3 cm., length without bill 52.5 cm., extent 78 cm.

No. 429 (coll. W. F. H.), ♀ March 8, 1901, the intermediate, measures: wing 25 cm., length without bill 53 cm., extent 87 cm.

No. 438 (coll. W. F. H.), March 27, 1901, ♂, typical *rubripes* measures: wing 28.5 cm., length without bill 55 cm., extent 94 cm.

Measurements taken in the flesh.

Mr. Brewster has given a very accurate description of this new subspecies, which is more common in Middle Southern Ohio in the spring than in the fall. All of these specimens were taken at Waverly, but I believe a careful investigation of the material in this state by Ohio ornithologists will prove *rubripes* to be found quite frequently almost anywhere.